

STUDIES ON FROGS OF THE GENUS *LEPTODACTYLUS*
(AMPHIBIA, LEPTODACTYLIDAE). V. TAXONOMIC
NOTES ON *L. LATINASUS*, *RHODONOTUS*,
ROMANI, AND *WUCHERRERI*

W. RONALD HEYER

Examination of type specimens during an ongoing biosystematic study of the frog genus *Leptodactylus* indicates the need for nomenclatural changes. The larva of one of the involved species is described and illustrated for the first time.

MATERIALS AND METHODS

The nomenclatural changes indicated herein resulted from examination of the types involved. I examined all the types personally, with the exception of *Cystignathus rhodonotus*. The methods for examining the adults and larvae are the same as those given earlier (Heyer, in press).

ACKNOWLEDGMENTS

The following curators have loaned material for examination (Museum abbreviations as used in text in parentheses): Javier Castroviño, Instituto "José de Acosta" de Zoología, Museo Nacional de Ciencias Naturales, Madrid (Madrid); B. Hubendick, Naturhistoriska Museet, Göteborg (Göteborg); Ernest E. Williams, Museum of Comparative Zoology, Harvard University (MCZ); and John W. Wright, Los Angeles County Museum of Natural History (LACM). Richard E. Etheridge, San Diego State College, examined the type of *C. rhodonotus* for me at the British Museum (BMNH). I thank him and David B. Wake, University of Chicago, for allowing me to report on the larvae which they collected. Jay M. Savage, University of Southern California, read and commented on the manuscript. The research and report were completed while I was a National Science Foundation Graduate Trainee at the University of Southern California.

SPECIES ACCOUNTS

Leptodactylus latinus

Leptodactylus latinus was described on the basis of a single specimen from Montevideo, Uruguay, in 1875 by Jiménez de la

