

# Notes on the frog fauna of the Amazon Basin

W. Ronald Heyer (\*)

## Abstract

The natural history of the frog fauna of the Amazon Basin is considered, based on data obtained from four major collections. Population structures in both forest and open environments are compared. Associations of species pairs in open formations are studied. It is estimated that at least 100 species are present in the Amazon Basin. Five speciation models are proposed to explain the high diversity of species.

I have had the opportunity to participate in two expeditions to the western Amazon region. This report deals with the natural history observations made on these trips together with an ecological analysis of the collections from these trips and collections from two other expeditions. All natural history observations and analyses on frogs are included here, rather than fragment the information. Taxonomic matters will be discussed in another paper.

The four expeditions on which this report is based are summarized briefly as:

1) Porto Velho: Departamento de Zoologia da Secretaria da Agricultura (now Museu de Zoologia, Universidade de São Paulo (MZUSP) expedition to Porto Velho, Rondônia, Brasil, from 28 October to 6 November 1954. This collection is represented by 17 species and 4086 individuals.

2) Expedição Permanente da Amazônia (EPA), supported by MZUSP, Fundação de Amparo à Pesquisa do Estado de São Paulo and Museu Goeldi; in 1973 a collaborative venture with the Museum of Comparative Zoology, Harvard University (Russell A. Mittermeier), from 20 August to 5 December 1973 along major rivers from Belém to Leticia, Colombia. Russell Mittermeier purchased 8238 individuals of 40 species incidental to the main purpose of the expedition.

3) Joint MZUSP and Smithsonian (USNM) expedition on the rio Purus, Brasil, from 2 December 1974 to 19 January 1975. This collection contains 811 specimens of 58 species.

4) Joint EPA-MZUSP and USNM expedition on the rio Madeira, Brasil from 31 October to 19 December 1975. The collection is represented by 1232 specimens of 53 species.

The itineraries of these trips will be described in the taxonomic paper where they are more appropriate.

## ECOLOGICAL GENERALIZATIONS

*Faunal — Habitat Associations:* At any locality in the Amazon basin, two major types of habitats are evident: forest and open. Of course, neither is a clear cut category for they grade into one another smoothly or abruptly, depending more or less on the human activity of the region. Open habitats have always been present, particularly at the edges of rivers. For all of the specimens collected by scientific personnel on the Madeira and Purus expeditions, the major habitat type was recorded for each specimen collected. Categories were either: (1) forest, (2) modified forest, such as selective logging or old second growth, and (3) open formations, such as river edges and human clearings. For purposes of analysis, any intermediate habitat, recorded in the field notes, was scored as the more forested category. The results of percent of individuals taken from each habitat type for the common species are presented in Table 1. The criterion used for common species in this case is 10 or more individuals (pooled localities), except for *Adenomera hylaedactyla* and *Hyla boans* for which calling specimens (not collected) supplement the actual specimens in the col-

(\*) — Division of Reptiles and Amphibians. National Museum of Natural History. Smithsonian Institution. Washington, D.C. (U.S.A.).

lection. If the rather rigid guideline of 80% occurrence in a single category is used, all but 4 species can be categorized as forest associated species (9) or open formation associated species (20). The four remaining species are primarily forest associated (*Phyllobates pictus*), primarily open formation associated (*Hyla lanciformis* and *perviceps*) or distributed in both forest and open formations (*Sphaenorhynchus eurhostus*). The association of most Amazonian frogs with one or the other major habitat type suggests that the species have evolved in either forests or open habitats, not both. Only in one case for those species tabled is the particular part of the breeding habitat for which the species is adapted found in both forests and open formations. This breeding habitat for *Sphaenorhynchus eurhostus* is a vegetation covered pond, either grass or *Pistia*.

The numbers of forests and open habitat associated species (Table 1) are not an accurate reflection of the true percentages of forests and open habitat associated species in the Amazonian anuran fauna. The open formation associated species were better sampled, because most specimens were collected from breeding chorus aggregations in open habitat ponds. In contrast, most forest associated species were collected individually from transects taken through the forest. This forest cruising technique produces few specimens but yields a high diversity. The distributions (Table 1) indicate that even with few habitat data, habitat association is reasonably accurate. Including the species with fewer than 10 individuals collected, there are presumable 38 forest associated species, 33 open formation associated species, and 2 species associated with both habitat types. As discussed below, the 73 species collected on the Madeira and Purus expeditions do not represent the total diversity of the Hylaeon anuran fauna. As the open formation fauna is easier to collect, I assume that most of the Hylaeon species not collected on the expeditions will be forest associated species.

All diurnal frogs are forest floor leaf litter species. All open formation associated frogs are nocturnal species. Most forest litter frogs

share the same type of nocturnal resting sites, i.e. leaves of herbaceous plants from 5 to 100 cm above the forest floor (*Bufo typhonius*, *Dendrophryniscus minutus*, both common *Phyllobates*, *Physalaenus petersi*). No *Adenomera andreae* were taken on leaves at night. Additional members of the forest litter fauna include *Dendrobates quinquevittatus*, *Phyllobates trivittatus*, and perhaps *Edalorhina* and one species of *Eleutherodactylus*. An approximation is that the forest litter frog fauna comprises about one quarter of the entire forest associated frog fauna.

TABLE 1. Distribution of individuals (in percentages) of common species among major habitat types. A = forest, B = modified forest, = open formation. Also see text.

SPECIES	A	B	C
<i>Adenomera andreae</i> .....	97	1	1
<i>A. hylaedactyla</i> .....	0	0	100
<i>Bufo granulosis</i> .....	0	0	100
<i>B. marinus</i> .....	4	2	93
<i>B. typhonius</i> .....	89	9	2
<i>Colostethus peruvianus</i> .....	86	14	0
<i>Dendrophryniscus minutus</i> .....	100	0	0
<i>Hyla</i> sp .....	100	0	0
<i>H. boans</i> .....	0	0	100
<i>H. fasciata</i> .....	84	4	11
<i>H. garbei</i> .....	83	17	0
<i>H. granosa</i> .....	8	0	92
<i>H. lanciformis</i> .....	26	6	68
<i>H. leali</i> .....	0	3	97
<i>H. leucophyllata</i> .....	11	4	86
<i>H. parviceps</i> .....	11	11	78
<i>H. punctata</i> .....	0	0	100
<i>H. raniceps</i> .....	0	2	98
<i>H. hodopepla</i> .....	6	0	94
<i>H. riveroi</i> .....	0	0	100
<i>H. rubra</i> .....	1	2	97
<i>H. triangulum</i> .....	0	0	100
<i>H. waldorfi</i> .....	0	2	98
<i>Leptodactylus bolivianus</i> .....	12	2	86
<i>L. fuscus</i> .....	0	6	93
<i>L. ocellatus</i> .....	0	0	100
<i>L. wagneri</i> .....	13	5	81
<i>Lysapsus limellus</i> .....	0	0	100
<i>Phrynohyas venulosa</i> .....	0	0	100
<i>Phyllobates femoralis</i> .....	91	9	0
<i>P. pictus</i> .....	72	28	0
<i>Physalaenus petersi</i> .....	100	0	0
<i>Sphaenorhynchus eurhostus</i> .....	45	0	55

*Species — Breeding Pond Associations:*  
If species share the same types of breeding ponds, the species assemblage probably evolved in the same environment.

Few forest ponds were found with breeding frogs. In those few forest ponds with breeding choruses, the species present are <sup>rare</sup> in museum collections. Many forest ponds, which I assumed to be ideal for anuran reproduction lacked any breeding frogs. Forest ponds may be used either for a brief time or be largely unsatisfactory. Both possible explanations may result from fish predation. Many ponds which are isolated at the beginning of the rainy season may become flooded and connected with river backflows. It is these flooded areas where many river fish also breed and would likely feed on any anuran larvae in the water. Also, erythrinid fishes occur in isolated ponds and creeks (Vanzolini, pers. comm.) Thus, many of the seemingly ideal breeding pools contain carnivorous fishes. One evolutionary response by the forest associated pond breeders would be to reproduce as soon as ponds develop and to have a brief larval life.

Data were recorded on species breeding in 28 open habitat ponds. The association between each species pair was evaluated by computing a coefficient of association, V, where

$$V = \frac{ad - bc}{(mnr)s} \sqrt{5}$$

where the right hand terms come from a 2X2 contingency table set up as

		Species B		
		present	absent	
Species A	present	a	b	m = a + b
	absent	c	d	n = c + d
		r = a + c	s = b + d	

where a, b, c, d are observed frequencies of species at breeding ponds (Pielou, 1969). V ranges from -1 (maximum negative association) to +1 (maximum positive association). Zero is no association. For the analysis herein, association values were calculated for only

those species recorded from 5 or more ponds. The results are presented in Table 2. Collins (1975) reported V values for breeding associations in a series of ponds in Michigan. He found that after applying different statistical tests to the contingency table, V values greater than .3 represented statistically valid association. If that criterion is used on the species data in Table 2, there are only 2 positive correlations: *S. eurhostus* — *H. leucophyllata* and *H. leali* — *L. wagneri*. This represents only 7% of the possible associations. In contrast, 25% of the temperate zone associations (Collins, 1975) were strongly positive. This difference suggests that the diversity of breeding ponds is lower and the available breeding period is shorter in Michigan than in Amazonia, a not unexpected explanation.

For the strongly negative associations (— .3) in Table 2, there are 5 pairs, but only one of them, *H. leali* — *H. riveroi*, involves closely related members.

The absence of positive and presence of negative associations that characterize Table 2 are consistent with a recent study that concluded that interspecific competition is evolutionarily unimportant in tadpole communities (Heyer, in press). One would expect evolution to act on biological interactions only where positive associations between species occur.

TABLE 2. Associations between the common open habitat species. Number in parentheses is the number of breeding ponds where the species was present.

	<i>H. lanciformis</i>	<i>H. leali</i>	<i>H. leucophyllata</i>	<i>H. riveroi</i>	<i>H. rubra</i>	<i>H. waldorfi</i>	<i>L. wagneri</i>	<i>S. eurhostus</i>
<i>H. lanciformis</i>	(7) 0.0	-.16	-.12	.22	-.03	.04	-.33	
<i>H. leali</i>		(8) .13	-.36	0.0	.11	.36	0.0	
<i>H. leucophyllata</i>			(8) .19	-.46	-.07	0.0	.55	
<i>H. riveroi</i>				(7) -.07	-.27	.33	.25	
<i>H. rubra</i>					(14) -.43	.07	-.25	
<i>H. waldorfi</i>						(5) .15	-.27	
<i>L. wagneri</i>							(7) .25	
<i>S. eurhostus</i>								(7)

It appears that the open formation fauna does not have a tightly integrated community structure, assuming breeding habitats are a reflection of the total community organization of frog communities.

**River Edge Frogs:** The following species are characteristic of the grassy river banks: *Adenomera hylaedactyla*, *Bufo granulosus*, *Hyla boans*, *Hyla granosa*, *Hyla punctata*, *Hyla raniceps* and *Leptodactylus fuscus*. Of these, *B. granulosus*, *H. punctata*, *H. raniceps*, and *L. fuscus* also occur in other open habitats away from the river edge. Apparently only *Hyla boans* and *Hyla granosa* breed in the river itself. Of the several fish seine samples taken in the Madeira and Purus rivers tadpoles were taken only once. As argued elsewhere (Heyer, McDiarmid, and Weigmann, 1975), the river tadpoles must have specific adaptations that allow them to occur with predatory fish.

**Adult Frog Food Resources:** One of the perplexing problems in analyzing resource utilization by adult frogs is the difficulty of gathering information on non-breeding individuals. With such data, one might examine ecological problems in frogs in the same successful manner as West Indian anoles have been studied. One approach appears to be a potentially fruitful one, and very preliminary results are offered in hopes that others might find the problem worth further study. When one searches the forest by night with flashlight, one finds frogs sitting on leaves, as well as arthropods sitting on leaves. One night, M. Heyer and I attempted to capture every arthropod and frog from leaf surfaces within reach for a one hour time period. Two frogs and 159 arthropods (Table 3) were collected. *Hyla lanciformis* had a 5 mm beetle and 25 mm preying mantis in the gut and *Hyla sp.* had a 7 mm grasshopper in the gut. The food items did not appear to be the same species as found in the general collections, but were comprised of the same kinds of potential prey items we collected. We collected about 80 arthropods per hour per person — it superficially appears as though food resources are superabundant in the forest for frogs. No palatability, catchability or digestive studies have been done for tropical forest frogs, however, so it

is not known what the true availability of food resources is for the frogs. A reasonable assumption at this point is that the nocturnal frogs seen on leaves at night in the forest are feeding on the arthropods also found on leaves at night (excluding diurnally active frogs which are resting on leaves at night, see above). Both the frogs and the arthropods are samplable resources.

TABLE 3. Arthropods collected on leaves at night in the forest.

Group	Size (classes) in mm	Number
Spiders	2	5
	3-4	2
	10	1
Cockroaches	6-10	2
	12-27	11
Beetles	3-4	4
	6-10	16
	10-20	11
Larval lepidopterans	10-12	2
	20-35	3
Homopterans	3	2
Ants	3	1
	11	1
Hemipterans	8	1
Dipterans	5	2
	13	1
	20	1
Moths	7-10	7
	16	1
Walking sticks	50-90	3
Orthopterans	4-7	3
	10-15	33
	15-20	24
	20-30	10
	30-40	8
	40-50	4

#### ANALYSIS OF RANDOM COLLECTIONS

The collections from the Madeira and Purus expeditions are not truly random collections because sampling was biased against the most common species. For example, not all adult *Bufo marinus* seen were collected.

The Porto Velho — MZUSP and EPA — Mittermeier collections are as random in nature as frog collections can be. All specimens collected were preserved, all specimens

possible were collected. The Porto Velho — MZUSP expedition did not have a herpetologist along, and Mittermeier purchased frogs from local children, a proven technique for lizards (e.g. Vanzolini, 1974).

One practical question to raise is whether the purchase technique is an effective way to sample the frog fauna of Amazonia. A comparison of the Porto Velho — MZUSP and EPA — Mittermeier collections with the Madeira and Purus collections show the former two collections have many more specimens than that of the Purus — Madeira collections; however, the latter two have a greater species diversity. The major discrepancy occurs with the forest associated fauna. Less than 0.01% of the specimens from the Porto Velho — MZUSP collection are forest associated species. No more than 5% of the EPA — Mittermeier specimens are forest associated species; these came from about half of the localities where Mittermeier purchased specimens. If the diurnal forest litter species are excluded, then less than 1% of the EPA — Mittermeier specimens are forest associated species from about one quarter of the localities. The Porto Velho — MZUSP and EPA — Mittermeier collections are mostly represented by open formation associated species. The purchasing technique in Amazonia, thus, is good for large samples of open formation associated species, weak for sampling forest associated species.

The most common (> 100 specimens) randomly collected open formation associated species can be analyzed in terms of relative abundance among collecting sites (Table 4). It is immediately obvious that what is common at one site is rare at another. Of the 280 *Leptodactylus ocellatus* collected, 254 came from a single site, Coari: 125 on 19 September 1973, 129 on 16-17 October 1973. Within a single season, it would appear that common species remain common. Of all the common species, *Bufo marinus* is the only one to approach a ubiquitous pattern of abundance and distribution.

This phenomenon of local rarity or abundance of species holds true over very small distances. At Borba, one hillside had many

forest floor frogs, the hillside opposite a small stream had almost none. At Puruzinho, we transected large areas of forest floor and found one very small area where we collected many more specimens than anywhere else, yet all the forest floor appeared similar.

TABLE 4. Common open habitat associated species in the EPA-Mittermeier collection.

Species	Total number of individuals	Greatest number of specimens from any single location-time period	% occurrence among all localities sampled
<i>B. marinus</i>	209	50	66
<i>H. geographica</i>	1205	1185	8
<i>H. punctata</i>	218	82	26
<i>H. rubra</i>	367	219	60
<i>L. fuscus</i>	164	146	8
<i>L. ocellatus</i>	280	129	13
<i>L. limellus</i>	5046	4737	18

#### ANURAN DIVERSITY

Two general questions concerning the Hylaeian fauna are: (1) Is the fauna uniformly distributed throughout the basin? and (2) Is the Andean Amazonian slope fauna more diverse than the basin fauna?

The Amazonian Andean slope frog fauna is among the most diverse faunas in the world. One site, Santa Cecilia, Ecuador, has been intensively sampled by field parties from the University of Kansas (e.g., Crump, 1974). The total of 81 species of frogs reported from Santa Cecilia is probably close to the true number of species present there. I worked at Limoncocha, Ecuador for two months, making collections of amphibians in the same way the Purus and Madeira collections were made. The known frog faunas of Limoncocha and Santa Cecilia are identical. Collecting curves were constructed for the Limoncocha, Purus, and Madeira collections, by plotting the number of frog species added per collecting day (Fig. 1).




Figure 1. Collecting effort curves for three Amazonian collections.

The collecting curves for the Limoncocha, Madeira, and Purus collections all start out similarly, but the Purus and Madeira curves are visibly steeper than the Limoncocha curve, indicating that the Madeira and Purus frog faunas contain more species than the Limoncocha fauna. Many of the species of the Madeira and Purus collections are the same, but there are several species represented in one collection and not the other. Most of these differences are due to seasonal differences in collecting times. The Madeira

collection was made in the transition period between the dry and wet season. Species such as *Bufo granulatus* and *Leptodactylus fuscus*, which breed at the very beginning of the wet season are well represented in the Madeira collection and absent in the Purus collections. Alternatively, several forest breeding hylids are represented in the Purus collections that are not present in the Madeira collections. The Madeira and Purus collections, back to back, span the transition from dry season to wet season, the greatest period of anuran breeding

TABLE 5. Preston canonical estimates (see text).

Collections	Number of species in collection	Predicted number of species in fauna
Limoncocha .....	52	72
Purus .....	58	87
Madeira .....	53	98
Purus + Madeira (P+M) ..	73	118
Porto Velho-MZUSP, EPA-Mittermeier, P+M, A .	81	150
Porto Velho-MZUSP, EPA-Mittermeier, P+M, B	81	109

activity. The Purus collection data, added to the Madeira data, forms a smooth collecting curve (Fig. 1) also indicating the two samples are from the same fauna. There is also the first indication of a leveling off at the end of this combined curve.

Preston (1962) proposed a formula for predicting the actual number of species based on a random sample of the fauna. As already discussed, the collections are not truly random, so any predictions will be approximate at best. The Preston canonical estimate is:

$$N = y_0 \cdot 6\sqrt{2\pi}, \text{ where}$$

$N$  = the predicted number of species,  $\delta$  = log standard deviation (Preston, 1962, p. 203 provides a table for this value based on the actual number of species in the collection), and  $y_0$  = the number of species in the modal octave, where the octaves are set up by ranking the numbers of individuals of each species collected. The estimated results for the collections are presented in Table 5. The one estimate for which the probable value is known (Limoncocha) is underestimated. The last two values of the table are estimated from combined data of the Porto Velho — MZUSP, EPA — Mittermeier, Madeira and Purus collections. The first estimate is using the modal octave, which in this case is the first octave, containing species represented by one or two specimens. The second estimate is based on the octave containing the next highest number of species and is included because it is more consistent with the other estimates.

At the moment, it would appear that a reasonable estimate of the number of species

in the Hylaeen frog fauna is at least 100. This is higher than the diversity at Santa Cecilia, but the geographic extent of the Hylaea is obviously much greater than Santa Cecilia. As is clear from Figure 1, collecting data for at least 40 days are needed from a single site in the basin proper to determine whether the diversity at a basin site is comparable to the slope fauna diversity.

#### ZOOGEOGRAPHY

The anuran fauna at Santa Cecilia is to date the most diverse studied (Crump, 1974). Recent revisionary studies have suggested that certain species have distributions associated with the upper Amazon basin of Colombia, Ecuador and Peru (e.g. Duellman, 1973, 1974a, b, Duellman and Crump, 1974). Groups of species with such distribution patterns would add to the overall diversity of the upper Amazon basin, contrasting with the presumed lower diversity of the lowland Amazon basin.

The four collections reported here are from the western half of the Brazilian Amazonian lowlands. Duellman and his associates (Duellman, 1973, 1974a, b, Duellman and Crump, 1974) have recently published studies based on previously available museum materials. In those studies, the following species were not known from the lowland Amazon basin and characterized as having upper Amazon distribution patterns: *Hyla fasciata*, *H. parviceps*, *H. sarayacuensis*, *Phyllomedusa palliata*, *P. tarsi*, and *Sphaenorhynchus carneus*. All of these species are represented in the collections from the western half of the Brazilian Amazonian lowlands. It is obviously premature to characterize individual distribution patterns as upper Amazonian. However, when one compares the total species lists of Santa Cecilia against the 4 collections of this report, the differences are appreciable. About 49% of the species recorded from Santa Cecilia are not found in the lowland Amazon collections. About 45% of the species in the lowland Amazon collections are not recorded from Santa Cecilia. Some of these differences are accounted for by differences in taxonomic opinion. These amount to no more than 7%

